arXiv:1601.03070v1 [hep-ph] 12 Jan 2016

Available online at www.sciencedirect.com
ScienceDirect
Nuclear Physics A 00 (2016) 1–4
Nuclear Physics A
www.elsevier.com/locate/procedia

The XXVth International Conference on Ultrarelativistic Nucleus-Nucleus Collisions

Direct photon production and jet energy-loss in small systems

Chun Shen[a], Chanwook Park[a], Jean-François Paquet[a,b], Gabriel S. Denicol[a,c], Sangyong Jeon[a], Charles Gale[1a]

[a]*Department of Physics, McGill University, 3600 University Street, Montreal, QC, H3A 2T8, Canada*
[b]*Department of Physics & Astronomy, Stony Brook University, Stony Brook, NY 11733, USA*
[c]*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

Abstract

Two types of penetrating probes, direct photon and QCD jets, are investigated in the background of a small and rapidly expanding droplet of quark-gluon plasma. The additional thermal electromagnetic radiation results in a ∼50% enhancement of the direct photons. In high multiplicity p+Pb collisions, jets can lose a sizeable fraction of their initial energy, leading to a charged hadron $R_{\rm pA}$ of ∼0.8 at a transverse momentum around 10 GeV. Those two proposed measurements can help understand the apparent collective behaviour observed in small collision systems.

Keywords: Direct photons, jet energy loss, small collision systems

1. Introduction

Signatures usually associated with hydrodynamic behaviour have been recently observed in high and intermediate multiplicity proton-nucleus (p+Pb) collisions at the Large Hadron Collider [1, 2, 3]. Even though those measurements are suggestive of the creation of a strongly coupled quark-gluon plasma (QGP), additional confirmation involving complementary observables needs to be pursued.

In this contribution we calculate the thermal photon radiation produced by a small and rapidly expanding QGP droplet, and we evaluate the energy loss of high energy QCD jets. We find that a

[1]Speaker.

significant amount of electromagnetic radiation is emitted, with thermal photons accounting for ∼50% of the direct photons produced in high multiplicity p+Pb collisions at low p_T. Furthermore, we show that in spite of the small system size, jets still lose a sizeable fraction of their initial energy, leading to a charged hadron $R_{\rm AA}$ of 0.8 at a transverse momentum of ∼10 GeV. Should these two measurements be within reach of the current generation of experiments, they would serve as additional evidence that a strongly coupled QGP is being produced in proton-nucleus collisions at the LHC. To complete the analysis, we also study direct photon production in other small systems, such as (p, d ^{3}He)+Au collisions at top RHIC energy.

2. Direct photons

In Pb+Pb collisions at $\sqrt{s_{\rm NN}}$ = 2.76 TeV, the measured direct photon yield shows a large enhancement over the photon spectra measured in pp collisions scaled by $N_{\rm coll}$, the number of binary collisions [4, 5]. This is usually quantified by the nuclear modification factor $R_{\rm AA}(p_T) \equiv \frac{dN^{\rm AA}}{dydp_T}/(N_{\rm coll}\frac{dN^{\rm pp}}{dydp_T})$. Direct photon $R_{\rm AA}$ in Pb+Pb collisions are shown in Figs. 1a-b, where it is seen that most of this enhancement - if not all - can be attributed to thermal radiation from the bulk medium [6]. Similarly, if a smaller medium achieves near-equilibrium, it will radiate thermal photons during its evolution. We investigated the thermal photon radiation from p+Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV and (p, d, ^{3}He)+Au at $\sqrt{s_{\rm NN}}$ = 200 GeV using the publicly available `iEBE-VISHNU` framework [7, 8]. Event-by-event hydrodynamic simulations with fluctuating Monte-Carlo Glauber initial conditions were used. Additional collision-by-collision fluctuations were considered in the initial energy density profiles, which led to a better description of charge hadron multiplicity distributions for these small collision systems [7]. The collision centrality was determined by the initial total entropy, dS/dy, which serves as a good approximation of the charged hadron multiplicity [9].



Fig. 1. *Panel (a)-(b):* Direct photon $R^{\gamma}_{\rm AA}$ in Pb+Pb collisions at LHC energy. *Panel (c)-(f):* The nuclear modification factor of direct photons in central and minimum bias collisions in small systems. *Panel (g)-(j):* The anisotropic flow coefficients $v_{2,3}$SP(scalar-product method) for direct photons in 0-1% and 0-20% centrality bins.

In Figs. 1c-f, we consider the direct photon signals below p_T < 3 GeV, and find that the thermal enhancement was about 50% in minimum bias collisions, and increased up to ∼150% for collisions in the 0-1% centrality class. Even though the space-time volumes of the proton-induced collisions are 20 times smaller than those in Pb+Pb collisions, the thermal signal is

only smaller by about a factor of 2. The direct photon measurements in minimum bias d+Au collisions still contain substantial statistical and systematic uncertainties[10]. Currently, none of the options - with or without thermal enhancement - are excluded by the data. The direct photon R_{AA}^{γ} increase as a function of p_T is faster for central collisions in p+Pb at $\sqrt{s_{\rm NN}} = 5.02\,$TeV and p+Au at $\sqrt{s_{\rm NN}} = 200\,$GeV than in (d, ^{3}He)+Au collisions at the top RHIC energy, owing to the fact that the number of fluctuating sources is smaller for p+A than for (d, ^{3}He)+A. Event-by-event fluctuations are more pronounced and will lead to higher temperatures in central p+A collisions. By correlating direct photon with charged hadrons, the direct photon anisotropic flow coefficients, $v_{2,3}\{\rm SP\}$, in 0-1% and 0-20% centrality bins are shown in Fig. 1g-j. The direct photon anisotropic flow in p+Pb collisions at 5.02 TeV is the largest among the four systems considered, because of larger pressure gradients and longer fireball lifetime.

3. Jet energy loss

Unlike direct photons that penetrate the fireball unscathed, high-energy partonic jets are strongly interacting probes. They lose some of their energy passing through the QGP. We estimate the amount of jet energy-loss in p+Pb collisions at $\sqrt{s_{\rm NN}} = 5.02\,$TeV using the Monte-Carlo event generator `MARTINI` [11]. Initial jets were generated using `Pythia` [12] with the standard tune settings [13]. Partons inside the proton were sampled from the parton distribution function (PDF), CTEQ6 [14]. For the Pb nucleus, additional cold nuclear effects were included by using EPS09 nuclear PDF [15]. These energetic partons were then evolved against the event-by-event hydrodynamic medium. Medium-induced radiation, elastic collisions, and path length dependence effects were taken into account in the model simulations [11, 16, 17]. Because the medium temperature is lower than perturbative scale, we choose a fixed strong coupling constant $\alpha_s = 0.27$ between jets and medium interaction. The strong coupling constant associated with medium-induced radiation is set to be running according to the jet momentum. All the partons were evolved to $T_c = 165\,$MeV and then hadronized according to the string fragmentation scheme germane to `Pythia`.


0-100% p+Pb @ 5.02 A TeV
pPb cold nuclear effect (EPS09)
pPb ebe $\alpha_s = 0.27$
ALICE $|\eta_{\rm CM}| < 0.3$
$R_{\rm pA}$
p_T (GeV)
0-1% p+Pb @ 5.02 A TeV
pPb cold nuclear effect (EPS09)
pPb ebe $\alpha_s = 0.27$
pPb smooth $\alpha_s = 0.27$
$R_{\rm pA}$
p_T (GeV)

Fig. 2. The nuclear modification factor of charged hadron, $R_{\rm pA}$, in minimum bias and 0-1% centrality p+Pb collisions at $\sqrt{s_{\rm NN}} = 5.02\,$TeV. Results in minimum bias collisions are compared with the ALICE measurement [18].

In Fig. 2, charged hadron $R_{\rm pA}$ in minimum bias and 0-1% centrality p+Pb collisions are shown for $10 < p_T < 100\,$GeV. In minimum bias collisions, the effect of energy loss is found to be negligible for $p_T > 40\,$GeV; only a modest mount of suppression is observed for $10 < p_T < 40\,$GeV. Both calculations - with and without energy loss - are consistent with the current ALICE

data [18]. On the other hand, a sizeable jet energy loss in the 0-1% central collisions is shown in the right panel of Fig. 2. Charged hadron $R_{\rm pA}$ is suppressed to ∼0.8 for $p_T \sim 10 - 20\,\text{GeV}$, and increases to ~ 1 around $p_T = 100\,\text{GeV}$. We also find that energy loss calculations with a smooth event-averaged hydrodynamic medium yield results for charged hadron $R_{\rm pA}$ very similar to those obtained with more realistic event-by-event simulations. The effects of event-by-event fluctuations on charged hadron $R_{\rm pA}$ is limited, for p + Pb collisions at the LHC. This is because the size of the entire fireball in the event-averaged hydrodynamic simulation is comparable to the typical total size of the hot spots in a fluctuating medium for p+Pb collisions. The charged hadron $R_{\rm pA}$ does not have the resolution to make such a distinction.

Acknowledgement

This work was supported by the Natural Sciences and Engineering Research Council of Canada. Computations were made on the Guillimin supercomputer at McGill University, managed by Calcul Québec and Compute Canada.

References

[1] B. B. Abelev, et al., Long-range angular correlations of π, K and p in p-Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV, Phys. Lett. B726 (2013) 164–177. arXiv:1307.3237, doi:10.1016/j.physletb.2013.08.024.

[2] V. Khachatryan, et al., Evidence for Collective Multiparticle Correlations in p-Pb Collisions, Phys. Rev. Lett. 115 (1) (2015) 012301. arXiv:1502.05382, doi:10.1103/PhysRevLett.115.012301.

[3] G. Aad, et al., Measurement of long-range pseudorapidity correlations and azimuthal harmonics in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, Phys. Rev. C90 (4) (2014) 044906. arXiv:1409.1792.

[4] J. Adam, et al., Direct photon production in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV,arXiv:1509.07324.

[5] B. Sahlmueller, Direct photon measurement in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV with ALICE, 2015. arXiv:1512.04230.

[6] J.-F. Paquet, C. Shen, G. S. Denicol, M. Luzum, B. Schenke, S. Jeon, C. Gale, The production of photons in relativistic heavy-ion collisions, arXiv:1509.06738.

[7] C. Shen, Z. Qiu, H. Song, J. Bernhard, S. Bass, U. Heinz, The iEBE-VISHNU code package for relativistic heavy-ion collisions, arXiv:1409.8164.

[8] C. Shen, J. F. Paquet, G. S. Denicol, S. Jeon, C. Gale, Thermal photon radiation in high multiplicity p+Pb collisions at the Large Hadron Collider, arXiv:1504.07989.

[9] C. Shen, Z. Qiu, U. Heinz, Shape and flow fluctuations in ultracentral Pb + Pb collisions at the energies available at the CERN Large Hadron Collider, Phys. Rev. C92 (1) (2015) 014901. arXiv:1502.04636.

[10] A. Adare, et al., Direct photon production in d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C87 (2013) 054907. arXiv:1208.1234, doi:10.1103/PhysRevC.87.054907.

[11] B. Schenke, C. Gale, S. Jeon, MARTINI: An Event generator for relativistic heavy-ion collisions, Phys. Rev. C80 (2009) 054913. arXiv:0909.2037, doi:10.1103/PhysRevC.80.054913.

[12] T. Sjstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, P. Z. Skands, An Introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159–177. arXiv:1410.3012.

[13] P. Skands, S. Carrazza, J. Rojo, Tuning PYTHIA 8.1: the Monash 2013 Tune, Eur. Phys. J. C74 (8) (2014) 3024. arXiv:1404.5630, doi:10.1140/epjc/s10052-014-3024-y.

[14] P. M. Nadolsky, H.-L. Lai, Q.-H. Cao, J. Huston, J. Pumplin, D. Stump, W.-K. Tung, C. P. Yuan, Implications of CTEQ global analysis for collider observables, Phys. Rev. D78 (2008) 013004. arXiv:0802.0007.

[15] K. J. Eskola, H. Paukkunen, C. A. Salgado, EPS09: A New Generation of NLO and LO Nuclear Parton Distribution Functions, JHEP 04 (2009) 065. arXiv:0902.4154, doi:10.1088/1126-6708/2009/04/065.

[16] S. Jeon, G. D. Moore, Energy loss of leading partons in a thermal QCD medium, Phys. Rev. C71 (2005) 034901. arXiv:hep-ph/0309332, doi:10.1103/PhysRevC.71.034901.

[17] S. Caron-Huot, C. Gale, Finite-size effects on the radiative energy loss of a fast parton in hot and dense strongly interacting matter, Phys. Rev. C82 (2010) 064902. arXiv:1006.2379, doi:10.1103/PhysRevC.82.064902.

[18] B. Abelev, et al., Transverse momentum distribution and nuclear modification factor of charged particles in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 110 (8) (2013) 082302. arXiv:1210.4520.